arXiv:1402.6084v1 [nlin.SI] 25 Feb 2014

GEOMETRIC REDUCTIONS OF ABS EQUATIONS ON AN n-CUBE TO DISCRETE PAINLEVÉ SYSTEMS

N. JOSHI, N. NAKAZONO, AND Y. SHI

Abstract. In this paper, we show for the first time how to relate n-dimensional cubes on which ABS equations hold to the symmetry group of discrete Painlevé equations. We focus here on the reduction to the q-discrete third Painlevé equation, which is a dynamical system on a rational surface of type $A_5^{(1)}$ with affine Weyl symmetry group $\widetilde{\mathcal{W}}\big((A_2+A_1)^{(1)}\big)$. This reduction also extends to other discrete Painlevé equations.

1. Introduction

We present a geometric method to obtain discrete Painlevé equations from higher-dimensional integrable discrete systems. Geometrically, symmetry groups of discrete Painlevé equations are affine Weyl groups, orthogonal to the divisor class of their initial value space in the Picard lattice [14, 16]. Higher dimensional discrete integrable systems arise from an entirely different geometric point of view, namely as multi-dimensionally consistent systems embedded on a hypercube, often referred to as consistency around a cube [13, 1, 2]. Unlike previous studies which have performed reductions of such equations via methods suited to specific examples [12, 8, 10, 9, 7, 15], we give a more general geometric construction, which provides affine Weyl groups of discrete Painlevé equations.

Multi-dimensionally consistent equations, with copies of the same equation holding on each face of a 3-dimensional cube, were classified by Adler *et al* [1]. The results are called ABS equations. Boll [4] extended these results to asymmetric 3-dimensional cubes, where equations on different faces may differ [3, 6]. These results were further extended to 4-dimensional cubes with the exception of asymmetric systems that incorporate H^6-type equations [6]. We construct a 4-dimensionally consistent system that does incorporate H^6-type equations and show its relation to the $A_5^{(1)}$-surface q-Painlevé system. Below, for conciseness, we refer to the n-dimensional cube as the n-cube.

This work is motivated by our previous findings [11], where quad-equations were observed on what is called the ω-lattice, constructed from the τ-function theory of the $A_5^{(1)}$-surface q-Painlevé system. The present paper begins with a system of quad-equations and travels in the opposite direction to obtain the q-Painlevé system, by reducing a 4-cube.

This research was supported by an Australian Laureate Fellowship # FL 120100094 and grant # DP130100967 from the Australian Research Council.

The plan of the paper is as follows. In section 2, we construct an asymmetric system of consistent quad-equations[i] on a 4-cube by fitting eight 3-cubes in a self-consistent way. In section 3, the reductions of this system are given. We show how to obtain $A_5^{(1)}$-surface q-Painlevé system by imposing a (1,1,1)-periodic condition along a symmetric 3-cube inside of the asymmetric 4-cube. An example with different step-length is also given. The latter example shows that our approach also includes previous periodic-type reductions on a 2D lattice. Finally, the paper ends with a conclusion.

2. Equations on an asymmetric 4-cube

To construct a 4-cube, we need four lattice directions l, m, n, k and four lattice parameters $\alpha(l), \beta(m), \gamma(n), \lambda(k)$. Let the dependent variable be $x = x(l, m, n, k)$ and denote its shifts in each direction respectively by

$$\bar{x} = x_1 = x(l+1, m, n, k), \tag{2.1a}$$
$$\hat{x} = x_2 = x(l, m+1, n, k), \tag{2.1b}$$
$$\tilde{x} = x_3 = x(l, m, n+1, k), \tag{2.1c}$$
$$\mathring{x} = x_4 = x(l, m, n, k+1). \tag{2.1d}$$

2.1. Symmetric 3-cube $\mathbf{C}_{312}$.

We start with the $H3$ quad-equation (2.2a) on a face of the 4-cube given by the l, n directions, where $x = x(l, n)$, $x_1 = x(l+1, n)$, $x_3 = x(l, n+1)$, $x_{13} = x(l+1, n+1)$. We construct a symmetric system of equations (2.2a)-(2.2f) on a 3-cube by taking auto-Bäcklund transformations of equation (2.2a):

$$Q(x, x_1, x_3, x_{13}; \alpha, \gamma) := xx_3 + x_1x_{13} - \alpha/\gamma(xx_1 + x_3x_{13}) = 0, \tag{2.2a}$$
$$Q(x, x_1, x_2, x_{12} : \alpha, \beta) = 0, \tag{2.2b}$$
$$Q(x, x_3, x_2, x_{23} : \gamma, \beta) = 0, \tag{2.2c}$$
$$Q(x_3, x_{13}, x_{23}, x_{123} : \alpha, \beta) = 0, \tag{2.2d}$$
$$Q(x_1, x_{13}, x_{12}, x_{123} : \gamma, \beta) = 0, \tag{2.2e}$$
$$Q(x_2, x_{12}, x_{23}, x_{123} : \alpha, \gamma) = 0, \tag{2.2f}$$

where x_2 denotes the auto-Bäcklund transformation of x with Bäcklund parameter β. We denote this 3-cube by C_{312}. See Figure 2.1a.

In this system we have four initial values x, x_1, x_2, x_3; and three quad-equations (2.2a)–(2.2c), adjacent to the vertex x. Vertices x_{12}, x_{13}, x_{23} are evaluated using the initial conditions and equations (2.2a)-(2.2c). x_{123} can then be evaluated using equations (2.2d)–(2.2f). 3D consistency of the system means that x_{123} evaluated by these three equations coincide. This is true in n-dimensions, i.e., it is multi-dimensionally consistent and hence it can be embedded on an n-cube. It is this remarkable fact we are going to utilise in deriving discrete Painlevé equations as periodic reductions on an n-cube. However, note that multi-dimensional consistent systems are not limited to systems with the same equation on all the faces.

[i] We use the term quad-equation throughout the paper to describe partial difference equations that relate the values the solution on the vertices of a quadrilateral.



(A) Symmetric 3-cube C_{312} (B) Asymmetric 3-cube C_{134}

FIGURE 2.1. Two types of 3-cubes C_{312} and C_{134} with different arrangements of equations on faces

.

2.2. **Asymmetric 3-cube $\mathbf{C}_{134}$.** Now consider a non-auto Bäcklund transformation of (2.3a) giving rise to different equations on different pairs of faces on the 3-cube. Here, we use Equations from Boll's classification of asymmetric 3-cubes [5] ((3.29)–(3.30) with $\delta_2 = \delta_3 = 0$). The 3-cube contains two H^4 type (H3) equations on the "Q" faces and four H^6 type equations for the"A" and "C" faces.

$$Q(x, x_1, x_3, x_{13}; \alpha, \gamma) := xx_3 + x_1x_{13} - \alpha/\gamma(xx_1 + x_3x_{13}) = 0, \tag{2.3a}$$
$$A(x, x_3, x_4, x_{34}; \delta_1) = xx_4 + x_3x_{34} + \delta_1 xx_3 = 0, \tag{2.3b}$$
$$C(x, x_1, x_4, x_{14}; \delta_1\eta_1) = xx_4 + x_1x_{14} + \delta_1\eta_1 xx_1 = 0, \tag{2.3c}$$
$$A' = A(x_1, x_{13}, x_{14}, x_{134}; \delta_1) = x_1x_{14} + x_{13}x_{134} + \delta_1 x_1x_{13} = 0, \tag{2.3d}$$
$$C' = C(x_3, x_{13}, x_{34}, x_{134}; \delta_1\eta_1) = x_3x_{34} + x_{13}x_{134} + \delta_1\eta_1 x_3x_{13} = 0, \tag{2.3e}$$
$$Q(x_4, x_{14}, x_{34}, x_{134}; \alpha, \gamma) = 0, \tag{2.3f}$$

where x_4 denotes the non-auto Bäcklund transformation of x and $\eta_1 = \alpha/\gamma$. The parameter δ_1 initially is a function of l, m, n and k, to be specified by the consistency conditions as we construct the 4-cube. Equations (2.3a)–(2.3f) make up the six quad-equations consistent on an asymmetric 3-cube. We denote this 3-cube by C_{134}, see Figure 2.1b.

Note that in obtaining Equations (2.3d)–(2.3f) we have used

$$\bar{\delta}_1 = \delta_1, \quad \tilde{\delta}_1\frac{\alpha}{\tilde{\gamma}} = \delta_1\frac{\alpha}{\gamma} \;\Rightarrow\; \frac{\tilde{\delta}_1}{\delta_1} = \frac{\tilde{\gamma}}{\gamma}. \tag{2.4}$$

Each of the above constructions can be repeated to obtain two more 3-cubes labelled C_{214} and C_{324}. C_{214} is obtained in the same way as C_{134} by replacing subscript 3 with 2 everywhere, with the condition $\eta_2 = \gamma/\beta$. C_{324} is obtained from C_{214} by replacing subscript 1 with 3 everywhere, with the condition $\eta_3 = \alpha/\beta$. In obtaining these two 3-cubes, we have imposed the following conditions

$$\delta_2 = \delta_3 = \delta_1\frac{\beta}{\gamma}, \quad \hat{\delta}_1 = \delta_1. \tag{2.5}$$

Equations (2.4) and (2.5) imply that

$$\delta_1(n, k) = \gamma(n)K(k), \tag{2.6}$$

where K is an arbitrary function of k.

We extend the above construction to four dimensions in the following way. The cube C_{3124} is obtained by shifting C_{312} in the k direction, C_{1342} by shifting C_{134} in the m direction, C_{2143} by shifting C_{214} in the n direction and C_{3241} by shifting C_{324} in the l direction. In summary, there are four 3-cubes adjacent to $x(l,m,n,k)$, namely, C_{312}, C_{134}, C_{214} and C_{324} and four 3-cubes adjacent to x_{1234}, namely C_{3124}, C_{1342}, C_{2143} and C_{3241}. These four 3-cubes are all 3D consistent under the conditions (2.4) and (2.5). Hence, we have eight three dimensionally consistent 3-cubes fitted consistently in a 4-cube. See Figure 2.2a.

The result contains two Bäcklund transformations, i.e., Equations (2.2f)–(2.3f), of the equation on the bottom face of the 4-cube, namely Equation (2.2). The permutability of these two Bäcklund transformations provides a system of 24 quad-equations, which can be embedded consistently on the 24 faces of the 4-cube. By construction, the system of equations on this 4-cube is four dimensionally consistent [6].



(A) A 4-cube.

(B) The 3-cube C_{214} is shaded pink and its tetrahedron equation is on the red vertices. Equation (3.4) holds on the face shaded blue.

FIGURE 2.2. The colours red, blue, green and yellow denote edges that are labelled by the same respective colours in Figure 2.1.

3. A (1,1,1)-reduction on the asymmetric 4-cube

We apply a periodic condition on the asymmetric 4-cube

$$\hat{\tilde{\tilde{x}}} = -i\lambda x, \tag{3.1}$$

where λ is only a function of k and $\overset{\circ}{\lambda} = q\lambda$. We call this a (1,1,1)-reduction of the 4-cube. We have the following conditions on the lattice parameters in the l, m, n directions: $\bar{\alpha}/\alpha = \hat{\beta}/\beta = \tilde{\gamma}/\gamma = q$, which are solved by $\alpha(l) = e^{\alpha_1} q^l$, $\beta(m) = e^{\beta_1} q^m$, $\gamma(n) = e^{\gamma_1} q^n$.

The periodic condition (3.1) enables us to evaluate $x(l,m,n,k)$ at any point in the lattice defined by the three directions l, m, n, using the initial conditions x, x_1, x_2, x_3 and the quad equations (2.2a)-(2.2c). To show that the reduced 4-cube can be extended also in the k direction, i.e., we can embed the reduce 4-cube in a four dimensional lattice, we derive the

expression relating iteration of $x(l,m,n,k)$ in the k direction (i.e., x_4) with iterations in two of the l,m,n directions, (for example x_1 and x_2).

On the asymmetric 3-cube C_{214}, shaded pink in Figure (2.2b), we have six equations for its six faces:

$$Q(x,x_1,x_2,x_{12}:\alpha,\beta)=0, \tag{3.2a}$$

$$A(x,x_2,x_4,x_{24};\delta_3)=xx_4+x_2x_{24}+\delta_1\beta\gamma^{-1}xx_2=0, \tag{3.2b}$$

$$C(x,x_1,x_4,x_{14};\delta_3\eta_3)=xx_4+x_1x_{14}+\delta_1\alpha\gamma^{-1}xx_1=0, \tag{3.2c}$$

$$A'=A(x_1,x_{12},x_{14},x_{124};\delta_3)=x_1x_{14}+x_{12}x_{241}+\delta_1\beta\gamma^{-1}x_1x_{12}=0, \tag{3.2d}$$

$$C'=C(x_2,x_{12},x_{24},x_{124};\delta_3\eta_3)=x_2x_{24}+x_{12}x_{124}+\delta_1\alpha\gamma^{-1}x_2x_{12}=0, \tag{3.2e}$$

$$Q(x_4,x_{14},x_{24},x_{124};\alpha,\beta)=0. \tag{3.2f}$$

From (2.2b), (3.2b) and (3.2e), we find

$$\frac{x_{12}}{x}\,\frac{x_{24}}{x_{12}}\,\frac{x}{x_{24}}=1 \quad\Rightarrow\quad \frac{(\alpha x_1-\beta x_2)(x_{124}+\alpha K x_2)}{(\beta x_1-\alpha x_2)(x_4+\beta K x_2)}=1. \tag{3.3}$$

(This follows from the tetrahedron property [1] and by "summing" the three-leg forms of the equations on the faces adjacent to x_2.) Moreover, from the face shaded blue in Figure 2.2b on which holds the equation for x_{12}, x_{124}, x_{1234}, and x_{123} we have another expression for x_{124}:

$$x_{1234}=-\frac{x_{12}(x_{124}+\gamma K x_{123})}{x_{123}} \;\Rightarrow\; -iq\lambda x_4=\frac{x_{12}(x_{124}-i\lambda\gamma K x)}{i\lambda x}, \tag{3.4}$$

where we have used the periodic condition (3.1) in rewriting the last equation. Using (3.4) to eliminate x_{124} in (3.3), we finally have

$$x_4=\frac{\left((-\alpha^2+\beta^2)x_1x_2-ix\gamma\lambda(\alpha x_1-\beta x_2)\right)K}{\lambda(\beta x_1-\alpha x_2)}. \tag{3.5}$$

4. Relation to q-Painlevé equations

In this section, we identify q-Painlevé equations from the (1,1,1)-reduction of the 4-cube described in §3. Observing that the reduction collapses the 3-cubes C_{312} and C_{3124} in Figure 2.2b to two copies of a hexagon, which can be extended everywhere in the triangular lattice, we find the affine Weyl group $\widetilde{\mathcal{W}}\big(A_2^{(1)}\big)$. But the fourth direction relating the two 3-cubes provides us with an extra direction that leads to a lattice with Affine Weyl symmetry group $\widetilde{\mathcal{W}}\big((A_2+A_1)^{(1)}\big)$, which is the full symmetry group of the $A_5^{(1)}$-surface q-Painlevé equation. A sub-case of the (1,1,1)-reduction leads us to the symmetric version of this equation, which is often called the second q-discrete Painlevé equation (q-P$_{\rm II}$).

Define

$$x(l,m,n,k)=(-1)^{\frac{1}{2}(l+m+n+k)}\lambda^m\omega(l,m,n,k), \tag{4.1}$$

then the periodic condition $\hat{\tilde{x}}=-i\lambda x$ implies

$$\hat{\tilde{\omega}}=\omega. \tag{4.2}$$

On letting $K=\lambda^{-1}(q\lambda^2-1)$, $\beta\alpha^{-1}=a_1$, $\alpha\gamma^{-1}=q^{-1}a_0$ and $\gamma\beta^{-1}=a_2$, our q-periodically reduced system on the asymmetric 4-cube is exactly the ω-lattice constructed from the τ-function frame work of $A_5^{(1)}$-surface q-Painlevé

system [11]. Results from [11] are reproduced in Appendix A to show that iterations of ω in the l, m, n and k direction give rise to the affine Weyl group $\widetilde{\mathcal{W}}\big((A_2+A_1)^{(1)}\big)$.

Figure 4.1 shows the vertices of the 4-cube now relabeled in ω variables. The quad-equations on the faces adjacent to ω are:

$$\frac{\lambda\hat{\tilde{\omega}}}{\omega} = \frac{\alpha\bar{\omega} - \beta\lambda\hat{\omega}}{-\beta\bar{\omega} + \alpha\lambda\hat{\omega}} \tag{4.3a}$$

$$\frac{\bar{\tilde{\omega}}}{\omega} = \frac{\alpha\bar{\omega} - \gamma\tilde{\omega}}{-\gamma\bar{\omega} + \alpha\tilde{\omega}} \tag{4.3b}$$

$$\frac{\lambda\hat{\tilde{\omega}}}{\omega} = \frac{\beta\lambda\hat{\omega} - \gamma\tilde{\omega}}{-\gamma\lambda\hat{\omega} + \beta\tilde{\omega}} \tag{4.3c}$$

$$\frac{\mathring{\bar{\omega}}}{\omega} = \frac{\alpha K}{\beta} + \frac{\mathring{\omega}}{\bar{\omega}} \tag{4.4a}$$

$$\frac{\mathring{\hat{\omega}}}{\omega} = \frac{K}{q\lambda} + \frac{\mathring{\omega}}{q\lambda^2\hat{\omega}} \tag{4.4b}$$

$$\frac{\mathring{\tilde{\omega}}}{\omega} = \frac{K\gamma}{\beta} + \frac{\mathring{\omega}}{\tilde{\omega}} \tag{4.4c}$$

The expression relating $\circ, \bar{}$ and $\hat{}$ directions is given by transforming Equation (3.5) to ω variables:

$$\mathring{\omega} = \frac{\lambda\big((-\alpha^2+\beta^2)\bar{\omega}\hat{\omega} - \gamma\omega(\alpha\bar{\omega} - \beta\lambda\hat{\omega})\big)K}{\beta(-1+q\lambda^2)(\beta\bar{\omega} - \alpha\hat{\omega})}. \tag{4.5}$$



FIGURE 4.1. Figure 2.2 relabelled in ω coordinates.

Using the fact $\hat{\bar{\tilde{\omega}}} = \omega$, we see that the inner 3-cube in Figure 4.1 collapses to a hexagon, and similarly for the outer 3-cube using $\mathring{\hat{\bar{\tilde{\omega}}}} = \mathring{\omega}$. Iterations of these provide two copies of the triangular lattice drawn in Figure 4.2.

Every vertex on the lower triangular lattice in Figure 4.2 can be calculated from the four initial conditions ω, $\bar{\omega}$, $\hat{\omega}$, $\tilde{\omega}$, using the three quad equations (4.3a)–(4.3c). (Vertices $\hat{\omega}, \bar{\tilde{\omega}}$ and $\hat{\tilde{\omega}}$ are calculated using (4.3a), (4.3b) and (4.3c) respectively.) Equation (4.5) provides a link between the upper and lower triangular lattices. In this way, we find that the iteration in the $\circ$ direction (or k direction) provides layers of triangular lattices. See Figure (4.2).

Without the (1,1,1)-periodic condition, the $H3$ system cannot be embedded into the layered triangular lattice which has affine Weyl symmetry $\widetilde{\mathcal{W}}\big((A_2+A_1)^{(1)}\big)$, which we call here $(A_2+A_1)^{(1)}$ lattice. So the system provided above is a reduction of the $H3$ system on a 4-cube. To show that the reduced 4-cube system is indeed q-Painlevé system on $A_5^{(1)}$-surface, we



FIGURE 4.2. 2 Hexagons from the outer and inner 3-cubes on $(A_2 + A_1)^{(1)}$ lattice

define:

$$f = \frac{\bar{\omega}}{\hat{\bar{\omega}}}, \quad g = \lambda \frac{\hat{\bar{\omega}}}{\omega} \tag{4.6}$$

and find their shifts in the $\bar{}$ direction, i.e., $\bar{f} = \bar{\bar{\omega}}/\hat{\bar{\bar{\omega}}}$, $\bar{g} = \lambda \hat{\bar{\bar{\omega}}}/\bar{\omega}$. For $\hat{\bar{\bar{\omega}}}$, we use (4.3b), shifted one step in $\bar{}$ and one step in $\hat{}$, along with the periodic condition (4.2). For $\bar{\bar{\omega}}$, we shift (4.3a) one step in $\bar{}$. These provide

$$\hat{\bar{\bar{\omega}}} = \frac{q\alpha\omega\bar{\omega} + \gamma\omega\hat{\bar{\omega}}}{\gamma\bar{\omega} + q\alpha\hat{\omega}}, \quad \bar{\bar{\omega}} = \frac{\beta\lambda\bar{\omega}\hat{\bar{\omega}} + q\alpha\lambda^2\hat{\bar{\omega}}\hat{\bar{\bar{\omega}}}}{q\alpha\bar{\omega} + \beta\lambda\hat{\bar{\bar{\omega}}}}. \tag{4.7}$$

Thus we have

$$\bar{g} = \frac{\lambda^2(1+ft)}{fg(f+t)}, \quad \bar{f} = \frac{\lambda^2(1+a\bar{g}t)}{f\bar{g}(\bar{g}+at)}, \tag{4.8}$$

where we have let

$$t = q\alpha/\gamma, \quad a = \gamma/\beta \tag{4.9}$$

and t is the independent variable of the q-Painlevé equation and a is a parameter. The system (4.8) is the q-discrete third Painlevé equation (q-$\mathrm{P_{III}}$) [16].

The triangular lattices in Figure 4.2 also provide a direct way of constructing the Bäcklund transformations of system (4.8). For translation in the $\circ$ direction for g and f we have $\overset{\circ}{g} = q\lambda\hat{\overset{\circ}{\omega}}/\overset{\circ}{\omega}$ and $\overset{\circ}{f} = \bar{\overset{\circ}{\omega}}/\hat{\overset{\circ}{\omega}}$. For $\hat{\overset{\circ}{\omega}}$ we use the equation on the face containing $\hat{\bar{\bar{\omega}}}$, $\hat{\omega}$, $\hat{\overset{\circ}{\omega}}$, $\hat{\overset{\circ}{\bar{\bar{\omega}}}}$ and the periodic condition

(4.2). These lead to

$$\mathring{\hat{\omega}} = \omega\left(-\frac{K\gamma}{q\beta} + \frac{\mathring{\omega}}{\hat{\omega}}\right). \tag{4.10}$$

Using (4.5), (4.10), and (4.4a) we have:

$$\mathring{g} = \frac{q(fgq + aft + a)}{g(fgq + aft + aq\lambda^2)}, \quad \mathring{f} = \frac{q(fg + aft\lambda^2 + a\lambda^2)}{f(fgq + aft + a)}, \tag{4.11}$$

which describe the Bäcklund transformation of f and g.

4.1. **(2,1) staircase reduction.** We now explain how different types of staircases taken on a two-dimensional square lattice (previously studied in the literature [15]) are actually sub-cases of the geometric reduction on the n-cube. We demonstrate this by investigating the only sub-case of the reduction considered in the previous section.

Let $l = n$, $\alpha = \gamma$, (i.e., $\bar{}=\tilde{}$) then the (1,1,1)-periodicity $\bar{\hat{\tilde{\omega}}} = \omega$ becomes the (2,1)-periodicity

$$\bar{\bar{\hat{\omega}}} = \omega. \tag{4.12}$$

System (4.3a)-(4.4c) reduces to one equation, i.e. Equation (4.3a) with the condition $\bar{\tilde{\alpha}}/\alpha = \hat{\beta}/\beta = q^2$. Using the same definition (4.6) for f and g, and (4.9) for t and a, we find

$$\bar{g} = \frac{\lambda^2}{gf}, \quad \bar{f} = \frac{g(at\bar{g} + 1)}{\lambda(\bar{g} + at)}, \tag{4.13}$$

which can be rewritten as a single equation for g

$$\bar{\bar{g}} = \frac{\lambda^3(\bar{g} + at)}{\bar{g}g(at\bar{g} + 1)}. \tag{4.14}$$

The resulting equation is the symmetric version of q-$\mathrm{P_{III}}$ (4.8), usually referred to as q-$\mathrm{P_{II}}$.

5. Conclusion

In this paper, we provided a new method of geometric reduction that relates ABS equations to discrete Painlevé equations. We constructed a q-Painlevé equation from an asymmetric system based on the $H3$ equation on a 4-cube and also provided the reduction to its symmetric form as a subcase. Bäcklund transformations of the discrete Painlevé equation arise as a natural by-product of our construction. Obtaining other structures related to integrability, such as Lax pairs, is also possible and will be reported in a separate paper. An interesting future direction is to extend our method to other types of discrete Painlevé equations classified by Sakai [16]. This will be the subject of future work.

Appendix A. The generators of $\widetilde{\mathcal{W}}\big((A_2+A_1)^{(1)}\big)$ and the triangular lattice

The affine Weyl group $\widetilde{\mathcal{W}}\big((A_2+A_1)^{(1)}\big)$ is generated by $s_0, s_1, s_2, \pi, w_0, w_1, r$, which are transformations of parameters and variables that satisfy the fundamental relations

$$s_i{}^2 = (s_i s_{i+1})^3 = \pi^3 = 1, \quad \pi s_i = s_{i+1}\pi, \quad (i \in \mathbb{Z}/3\mathbb{Z}), \tag{A.1}$$

$$w_0{}^2 = w_1{}^2 = r^2 = 1, \quad r w_0 = w_1 r. \tag{A.2}$$

Here, the action of $\widetilde{W}(A_2^{(1)}) = \langle s_0, s_1, s_2, \pi\rangle$ and that of $\widetilde{W}(A_1^{(1)}) = \langle w_0, w_1, r\rangle$ commute. Full details of how to construct this affine Weyl group from the ω-lattice can be found in [11], with the identification $\omega = \omega_0$. Here we point out how this construction can be related to the triangular lattice shown in Figure 4.2.

Define the translations T_i $(i = 1, 2, 3, 4)$ by

$$T_1 = \pi s_2 s_1, \quad T_2 = \pi s_0 s_2, \quad T_3 = \pi s_1 s_0, \quad T_4 = r w_0, \tag{A.3}$$

where T_i $(i = 1, 2, 3)$ are translations of $\widetilde{W}(A_2^{(1)})$ and T_4 is a translation of $\widetilde{W}(A_1^{(1)})$. We connect these generators to the triangular lattice in Figure 4.2 by identifying

$$\begin{aligned}
&T_1 : (a_0, a_1, a_2, \lambda) \to (q a_0, q^{-1} a_1, a_2, \lambda) \quad &&\Leftrightarrow \quad {}^{-} : \ l \mapsto l + 1,\\
&T_2 : (a_0, a_1, a_2, \lambda) \to (a_0, q a_1, q^{-1} a_2, \lambda) \quad &&\Leftrightarrow \quad \hat{} : \ m \mapsto m + 1,\\
&T_3 : (a_0, a_1, a_2, \lambda) \to (q^{-1} a_0, a_1, q a_2, \lambda) \quad &&\Leftrightarrow \quad \tilde{} : \ n \mapsto n + 1,\\
&T_4 : (a_0, a_1, a_2, \lambda) \to (a_0, a_1, a_2, q\lambda) \quad &&\Leftrightarrow \quad \circ : \ k \mapsto k + 1.
\end{aligned}$$

The periodicity condition $\hat{\tilde{\bar{\omega}}} = \omega$ corresponds to the relation $T_1 T_2 T_3 = 1$ of $\widetilde{W}(A_2^{(1)})$.

The generators s_0, s_1, s_2 are illustrated in Figure A.1.



Figure A.1. Reflections of $\widetilde{W}(A_2^{(1)})$

References

[1] V. E. Adler, A. Bobenko, and Y. B. Suris. Classification of integrable equations on quad-graphs. The consistency approach. *Communications in Mathematical Physics*, 233(3):513–543, 2003.

[2] V. E. Adler, A. I. Bobenko, and Y. B. Suris. Discrete nonlinear hyperbolic equations. Classification of integrable cases. *Functional Analysis and its Applications*, 43(1):3–17, 2009.
[3] J. Atkinson. Bäcklund transformations for integrable lattice equations. *J. Phys. A*, 41(13):135202, 8, 2008.
[4] R. Boll. Classification of 3D consistent quad-equations. *J. Nonlinear Math. Phys.*, 18(3):337–365, 2011.
[5] R. Boll. Corrigendum: "Classification of 3D consistent quad-equations". *J. Nonlinear Math. Phys.*, 19(4):1292001–1–1292001–3, 2012.
[6] R. Boll. On Bianchi permutability of Bäcklund transformations for asymmetric quad-equations. *ArXiv e-prints*, Nov. 2012.
[7] C. M. Field, N. Joshi, and F. W. Nijhoff. q-difference equations of KdV type and Chazy-type second-degree difference equations. *J. Phys. A*, 41(13):332005, 13, 2008.
[8] B. Grammaticos, A. Ramani, J. Satsuma, R. Willox, and A. Carstea. Reductions of integrable lattices. *Journal of Nonlinear Mathematical Physics*, 12(suppl 1):363–371, 2005.
[9] M. Hay, J. Hietarinta, N. Joshi, and F. Nijhoff. A Lax pair for a lattice modified KdV equation, reductions to q-Painlevé equations and associated Lax pairs. *Journal of Physics A: Mathematical and Theoretical*, 40(2):F61, 2006.
[10] N. Joshi, B. Grammaticos, T. Tamizhmani, and A. Ramani. From integrable lattices to non-QRT mappings. *Letters in Mathematical Physics*, 78(1):27–37, 2006.
[11] N. Joshi, N. Nakazono, and Y. Shi. ABS equations in τ-functions framework of $A_5^{(1)}$-surface q-Painlevé systems. *ArXiv e-prints*, Jan. 2014.
[12] F. Nijhoff and V. Papageorgiou. Similarity reductions of integrable lattices and discrete analogues of the Painlevé II equation. *Physics Letters A*, 153(6):337–344, 1991.
[13] F. W. Nijhoff and A. J. Walker. The discrete and continuous Painlevé VI hierarchy and the Garnier systems. 2001.
[14] K. Okamoto. Sur les feuilletages associés aux équations du second ordre à points critiques fixes de P. Painlevé, espaces des conditions initiales. *Science Council of Japan. Japanese Journal of Mathematics. New Series*, 5:1–79, 1979.
[15] C. Ormerod. Reductions of lattice mKdV to q-PVI. 376(45):2855–2859, 2012.
[16] H. Sakai. Rational surfaces associated with affine root systems and geometry of the Painlevé equations. *Communications in Mathematical Physics*, 220(1):165–229, 2001.

School of Mathematics and Statistics F07, The University of Sydney, NSW 2006, Australia, Tel: +61 2 9351 2172, Fax: +61 2 9351 4534
E-mail address: nalini.joshi@sydney.edu.au

School of Mathematics and Statistics F07, The University of Sydney, NSW 2006, Australia, Fax: +61 2 9351 4534
E-mail address: nobua.n1222@gmail.com

School of Mathematics and Statistics F07, The University of Sydney, NSW 2006, Australia, Fax: +61 2 9351 4534
E-mail address: yshi7200@gmail.com